

# TURFGRASS SCIENCE

at the UT Institute of Agriculture

## White Clover (*Trifolium repens*)

Gregory K. Breeden, Extension Specialist, Turfgrass Weed Science

James T. Brosnan, Associate Professor, Turfgrass Weed Science

Department of Plant Sciences

### Introduction

White clover (*Trifolium repens*) is a perennial broadleaf weed found in both warm- and cool-season turf. There are numerous clover species, but white clover is the most common species found in turfgrass. White clover can persist at most mowing heights and in both areas of full sun to partial shade. A leguminous weed, white clover is commonly found in areas of low nitrogen fertility. Legumes are plants that have the ability to fix (i.e., obtain) nitrogen from the atmosphere; thus, white clover can persist in areas that have not been fertilized enough to sustain dense, vigorous turf growth. While not one of the more difficult-to-control broadleaf weeds in Tennessee, it is often perceived as one of the most persistent.

### White Clover Identification

White clover is a prostrate, spreading, perennial legume that spreads from stems and stolons that root at the nodes (Figure 1). It can be found in clusters or mats throughout the year, but grows most vigorously in spring and fall. White clover leaves are arranged in a trifoliate structure; three oval to elliptical-shaped leaflets comprise a single trifoliate leaf (Figure 2). Leaves often have a white watermark that partially encircles the base of each leaflet. Leaves are also slightly toothed along the margins. White clover produces circular white flowers (sometimes with a pink hue) on long stems that appear during the spring and fall (Figure 3).




Figure 1. White clover (*Trifolium repens*) in turf



Figure 2. White clover (*Trifolium repens*) leaves



Figure 3. White clover (*Trifolium repens*) flowers

## White Clover Control Options

### Cultural Practices

The best defense against any weed infestation is maintaining a dense, vigorous stand of turfgrass. Growing conditions that favor turf often discourage the growth of white clover. Increasing nitrogen fertility and reducing soil moisture will help create an environment better-suited for turfgrass growth and less conducive for white clover. Additionally, soil test regularly and apply nutrients according to soil test

recommendations. While cultural practices can be used to help discourage white clover infestations, they will not provide complete control. Herbicide applications will be required for complete eradication.

### Herbicide Options

There are no preemergence herbicide options for control of white clover. Numerous herbicide options exist for postemergence control of white clover. Herbicides that contain three or more phenoxy (or phenoxy-type) herbicides can be used to control white clover in established turf. Herbicides containing the active ingredients aminocyclopyrachlor, fluroxypyr, triclopyr or quinclorac have performed well in research at the University of Tennessee. Aminocyclopyrachlor can be found in the herbicide Imprelis. Fluroxypyr can be found in herbicides such as Spotlight (fluroxypyr) and Escalade II (fluroxypyr, 2,4-D and MCPP), while quinclorac can be found in herbicides such as Drive XLR8 (quinclorac) and Solitare (quinclorac and sulfentrazone). Most of these herbicides can be used on both warm- and cool-season turfgrasses grown throughout Tennessee.

**Table 1. Postemergence herbicides for white clover (*Trifolium repens*) control**

Product Name	Active Ingredients	Rate/ac	Turf Species
Blindside	metsulfuron + sulfentrazone	3.25-10 oz.	Kentucky Bluegrass, Tall Fescue, Bermudagrass, Zoysiagrass, Centipedegrass
Celsius	thiencarbazone + iodosulfuron + dicamba	2.5-3.75 oz.	Bermudagrass, Zoysiagrass, Centipedegrass
Confront	triclopyr + clopyralid	1-2 pts.	Kentucky Bluegrass, Fine Fescue, Tall Fescue, Perennial Ryegrass, Bermudagrass, Zoysiagrass, Centipedegrass
Drive XLR8	quinclorac	2 qts.	Kentucky Bluegrass, Fine Fescue, Tall Fescue, Perennial Ryegrass, Bermudagrass, Zoysiagrass
Escalade II	2,4-D + fluroxypyr + dicamba	2-3 pts.	Kentucky Bluegrass, Fine Fescue, Tall Fescue, Perennial Ryegrass, Bermudagrass, Zoysiagrass
Manor	metsulfuron	0.125-1 oz.	Kentucky Bluegrass, Fine Fescue, Bermudagrass, Zoysiagrass, Centipedegrass
Millennium Ultra 2	2,4-D + clopyralid + dicamba	2-3 pts.	Kentucky Bluegrass, Fine Fescue, Tall Fescue, Perennial Ryegrass, Bermudagrass, Zoysiagrass
Momentum fx2	2,4-D + triclopyr + fluroxypyr	3-4 pts.	Kentucky Bluegrass, Fine Fescue, Tall Fescue, Perennial Ryegrass, Bermudagrass, Zoysiagrass, Centipedegrass
Monument	trifloxysulfuron	0.53 oz.	Bermudagrass, Zoysiagrass
Powerzone	carfentrazone + MCPA + MCPP	2-4 pts. + dicamba	Kentucky Bluegrass, Fine Fescue, Tall Fescue, Perennial Ryegrass, Bermudagrass, Zoysiagrass
Pylex	topramezone	1-1.5 oz.	Kentucky Bluegrass, Fine Fescue, Tall Fescue, Perennial Ryegrass, Centipedegrass

**Table 1. Postemergence herbicides for white clover (*Trifolium repens*) control (continued)**

Product Name	Active Ingredients	Rate/ac	Turf Species
Q-4	quinclorac + sulfentrazone + 2,4-D + dicamba	7-8 pts.	Kentucky Bluegrass, Fine Fescue, Tall Fescue, Perennial Ryegrass, Bermudagrass, Zoysiagrass (Dormant)
Solitare	quinclorac + sulfentrazone	16-32 oz.	Kentucky Bluegrass, Fine Fescue, Tall Fescue, Perennial Ryegrass, Bermudagrass, Zoysiagrass, Centipedegrass
Speedzone	carfentrazone + 2,4-D + MCPP + dicamba	3-5 pts.	Kentucky Bluegrass, Fine Fescue, Tall Fescue, Perennial Ryegrass, Bermudagrass, Zoysiagrass
SquareOne	carfentrazone + quinclorac	8-18 oz.	Kentucky Bluegrass, Fine Fescue, Tall Fescue, Perennial Ryegrass, Bermudagrass, Zoysiagrass, Centipedegrass
Surge	sulfentrazone + 2,4-D + MCPP + dicamba	3-4 pts.	Kentucky Bluegrass, Fine Fescue, Tall Fescue, Perennial Ryegrass, Bermudagrass, Zoysiagrass
Three-way Selective, Trimec Classic, Trimec Southern, Triplet, Others	2,4-D + MCPP + dicamba	Product-Dependent	Product-Dependent
Tribute Total	thiencarbazono + foramsulfuron + halosulfuron	2-3.2 oz.	Bermudagrass, Zoysiagrass
Turflon Ester	triclopyr	0.5-1 qt.	Kentucky Bluegrass, Tall Fescue, Perennial Ryegrass

Many of the sulfonylurea herbicides also control white clover in warm-season turf, including chlorsulfuron (Corsair), flazasulfuron (Katana) and trifloxysulfuron (Monument). Metsulfuron can also be used for white clover control. Metsulfuron can be applied alone (Manor) and in combination with other herbicides such as sulfentrazone (Blindside). See Table 1 for a complete list of herbicides active against white clover. Regardless of the herbicide option chosen, follow-up applications may be needed for complete control.

### Final Thoughts

Incorporating improved cultural practices with timely herbicide applications will provide long-term white clover control. Multiple herbicides options exist for effective control of white clover in turf.

Always refer to the product label for specific information on proper product use, tank-mix compatibility and turfgrass tolerance. Mention of trade names or commercial products in this publication is solely for the purpose of providing specific information and does not imply recommendation or endorsement by the University of Tennessee Institute of Agriculture.

For more information on turfgrass weed control, visit the University of Tennessee's turfgrass weed science website, [tennesseeturfgrassweeds.org](http://tennesseeturfgrassweeds.org).

Herbicides listed in this publication have provided good to excellent control in research trials conducted at the University of Tennessee; however, other herbicides may also have activity on this weed. For more information on herbicide selection, please visit University of Tennessee Mobile Weed Manual (MWM) at [mobileweedmanual.com](http://mobileweedmanual.com). MWM was developed by UT Extension professionals to assist green industry professionals in selecting herbicides for use in turf and ornamentals. MWM is a web-based platform optimized for use on mobile devices such as smartphones and tablets, but it will function on desktop and laptop computers as well. The site provides users with weed control efficacy information for 90 different herbicides, tolerance information for over 2,300 turf and ornamental species, as well as direct links to label and material safety data sheet information on herbicides used for turf and ornamental weed management.



Disclaimer

This publication contains herbicide recommendations that are subject to change at any time. The recommendations in this publication are provided only as a guide. It is always the herbicide applicator's responsibility, by law, to read and follow all current label directions for the specific herbicide being used. The label always takes precedence over the recommendations found in this publication.

Use of trade or brand names in this publication is for clarity and information; it does not imply approval of the product to the exclusion of others that may be of similar, suitable composition, nor does it guarantee or warrant the standard of the product. The author(s), the University of Tennessee Institute of Agriculture and University of Tennessee Extension assume no liability resulting from the use of these recommendations.

[AG.TENNESSEE.EDU](http://AG.TENNESSEE.EDU)

Real. Life. Solutions.

W 267 02/15 (Rev.) 15-0034

Programs in agriculture and natural resources, 4-H youth development, family and consumer sciences, and resource development.  
University of Tennessee Institute of Agriculture, U.S. Department of Agriculture and county governments cooperating.  
UT Extension provides equal opportunities in programs and employment.